

Composition : Terbinafine 250 mg as Terbinafine Hydrochloride Tablet.

Indication : Mycofree tablet is indicated for the treatment of :

- Onychomycosis of fingernails and toenails due to dermatophytes.
- Fungal infections of the skin (Tinea corporis, Tinea cruris, Tinea pedis).
- Hair and scalp infections (Tinea capitis).
- Yeast infections of the skin caused by the genus Candida (e.g. Candida albicans), where oral therapy is generally considered appropriate owing to the site, severity or extent of the infection.

Dosage and administration :

For Tinea cruris: 250mg tablet once daily for 2 to 4 weeks.

For Tinea pedis: 250mg tablet once daily for 2 to 6 weeks.

For Tinea corporis: 250mg tablet once daily for 4 weeks.

For dermatophyte infections of the finger & toenails : 250mg tablet once daily for 6 to 12 weeks. Longer treatment is necessary for toenail infections.

For skin and hair infections : 250mg tablet once daily for 2 to 6 weeks.

Or, as directed by the registered physician.

Side effects : Side effects are generally mild to moderate and transient, most common side effects are: Gastrointestinal symptoms-dyspepsia, loss of appetite, nausea, mild abdominal pain, diarrhoea, headache and dizziness.

Contraindication : It is contraindicated in patients with known hypersensitivity to the drug.

Drug Interactions : Plasma

Mycofree Tablet

concentration of Terbinafine may be increased by drugs that inhibit its metabolism by cytochrome P450, such as Cimetidine and decreased by drugs such as Rifampicin that induce cytochrome P450.

Precaution : It should be used with caution in patients with impaired hepatic and renal function.

Pregnancy and lactation : Pregnancy : Pregnancy category B. It should not be used during pregnancy unless the potential benefits clearly outweigh the risk. **Lactation :** It should not be given to nursing mothers because the drug is excreted in breast milk.

Packing : Mycofree-250 : 1 x 10's tablets in blister pack.