

Composition : Dexamethasone 1.0 mg / ml.

Indications : Eye : Non-infectious forms of conjunctivitis, keratitis and blepharitis, especially of allergic origin. Inflammation of the anterior uvea (iritis, iridocyclitis), scleritis, episcleritis, orbital myositis and sympathetic ophthalmia.

Ear : Redness, itching and swelling caused by certain ear problems.

Dosage and administration : Eye : Administer 1 drop 3-5 times daily into the conjunctival sac.

Acute cases : up to 1 drop hourly.

Ear : 3 or 4 drops in the ear two or three times a day.

Or, as directed by the registered physician.

Contraindications : Hypersensitivity to the ingredients, injuries and ulcerous processes of the cornea, Herpes simplex and other viral infections, mycosis, glaucoma. Similarly, Celudex 0.1% should not be administered to newborn babies.

Precautions : If no improvement is observed after 7-8 days treatment, other therapeutic measures should be considered. Due to the risk of systemic response, the administration to children under 2, is not advisable. During treatment with Celudex 0.1% the intraocular pressure should be measured regularly. Patients wearing contact lenses must not use the eye drops during the time the lenses are worn.

Drug interaction : None relevant to topical use.

Celudex

Sterile Eye & Ear Drops

Side effects : After several weeks administration to predisposed patients there may be a reversible increase in intraocular pressure. Systemic adverse reactions may occur after long-term topical application of corticosteroids (above all in children). Isolated cases of cataract and corneal softening have been reported after long-term treatment with steroids.

Pregnancy and lactation : Pregnancy Category **C** by FDA.

Packing : 5 ml in plastic dropper bottle.