

Composition : Each ml contains Povidone BP 50 mg.

Indications : Povidone Eye Drops is used for the symptomatic treatment of dry eye conditions including Keratoconjunctivitis sicca. It is also given as a substitute of tear fluid in case of unstable tear film or insufficient moistening of the eye surface.

Dosage and administration : 1 drop 4 times daily or as required, depending upon the severity of the disease, to be instilled into the conjunctival sac. Or, as directed by the registered physician.

Contraindication : It is contraindicated in patients with known hypersensitivity to any ingredient of the product.

Use in Pregnancy and lactation : There is no experience regarding the safety of the Povidone Eye Drops in human pregnancy or lactation. Administration during pregnancy and lactation is therefore not recommended, except for compelling reasons.

Side effects : Occasionally mild-transient burning or sticky sensation and very rarely irritation or hypersensitivity reactions are reported. Blurred vision after application may occur.

Precautions : Patients who experience blurred vision after application of the product should not drive or use machinery until their vision has cleared. Contact lenses should not be worn during instillation of the product. After instillation there should be an interval of at least 30 minutes before reinsertion of contact lenses.

Drug interactions : If several medicines are to be administered to the eye, there

Eyetear

Sterile Eye Drops

should be an interval of at least 5 minutes between each application.

Overdose : Not known and not likely about overdose.

Storage : Store in a cool & dry place. Protected from light. Keep out of reach of children. Do not touch the dropper tip to surfaces since this may contaminate the solution. Do not use after 1 month of first opening.

Packing : 10 ml in plastic dropper bottle.