

Composition : Vitamin A 50,000 I.U. & 2,00,000 I.U. Soft Capsule.

Indications : Illness due to vitamin A deficiency in ophthalmology such as night blindness, xerophthalmia and dermatological such as changes in skin, hair and nails. Concomitant therapy of mucosa illnesses such as sinusitis, bronchitis, in acne vulgaris, ichthyosis, Darier's disease, psoriasis etc.

To meet vitamin A demand in growth, resistance to infections and night blindness. Ratinol Forte is also indicated to meet vitamin A deficiency after diarrhoea and prophylaxis of measles.

Dosages : For Adults : 50,000 I.U - 1,00,000 I.U daily up to 2,00,000 I.U if necessary.

Children :

| Disease | Age | Dosage | Duration of treatment |
|---|--------------|--------------|----------------------------|
| Night blindness Bitot's spots Xerophthalmia | Above 1 year | 2,00,000 I.U | 1st day, 2nd day, 14th day |
| Measles | Above 1 year | 2,00,000 I.U | 1st day, 2nd day |
| Diarrhoea Respiratory tract infection | Above 1 year | 2,00,000 I.U | Every time after disease |
| Severe malnutrition | Above 1 year | 2,00,000 I.U | Single dose |

or, as directed by the registered physician.

Side effects : Vitamin A intoxication includes irritability, vomiting, loss of appetite, headache, dry and pruritic skin, skin desquamation, fatigue, pain in ankles and feet, myalgia, loss of body hair, papilledema, nystagmus, liver

Ratinol Forte Capsule

sclerosis and cirrhosis.

Contraindication : Hypervitaminosis of vitamin A.

Precautions : Ensure Ratinol Forte free interval after long term therapy with vitamin A. No daily dose over 5,000 I.U. during pregnancy. Vitamin A doses over 50,000 I.U. under medical supervision only.

Packing :

Ratinol Forte-50,000 : 6 x 10's soft capsules blister pack.

Ratinol Forte-2 lac : 5 x 10's soft capsules blister pack.