

Composition : (Dexamethasone 1mg+ Neomycin Sulfate 3,500 IU+ Polymyxin B Sulfate 6,000 IU)/ml.

Indications : Neo-DP is indicated for the short term treatment of steroid responsive conditions of the eye when prophylactic antibiotic treatment is also required, after excluding the presence of fungal and viral disease.

Dosage & administration : Apply one or two drops to each affected eye up to six times daily or more frequently if required. Severe infections may require one or two drops every 15-20 minutes initially, reducing the frequency of instillation gradually as the infection is controlled. Or as directed by registered physician.

Contraindications : Hypersensitivity to any component of the preparation. Herpes simplex and other viral diseases of the cornea and conjunctiva. Fungal disease and ocular tuberculosis.

Side effects : Reactions occurring most often are allergic sensitisations. The reactions due to the steroid component are elevation of intraocular pressure (IOP) leading to optic nerve damage, posterior subcapsular cataract formation and delayed wound healing.

Precautions : Prolonged use should be avoided as it may lead to skin sensitisation and emergence of resistant organisms. Because of risk of steroid glaucoma and cataract formation, the

Neo-DP

Sterile Ophthalmic Suspension

intraocular pressure and lens must be checked frequently during use of the preparation.

Packing : 5ml in plastic dropper bottle.