

**Composition :** Diclofenac sodium  
1.00mg/ml.

**Indications :** Postoperative inflammation after cataract surgery and other surgical procedures. Preoperative and postoperative prevention of cystoid macular oedema associated with lens extraction and intraocular lens implantation. Non-infectious inflammatory conditions affecting the anterior region of the eye (e.g. chronic non-infectious conjunctivitis). Post-traumatic inflammation in penetrating wounds (as an adjuvant to local anti-infective therapy).

**Dosage and administration :**  
Preoperative : up to 5x1 drop over the 3 hours preceding surgery. Postoperative : 3x1 drop immediately after surgery, thereafter 3-5x1 drop daily for as long as required.

**Other indications :** 4-5x1 drop daily, depending on the severity of the condition. Or as directed by the physician.

**Contraindications :** Like other non-steroidal anti-inflammatory agents, Diclofenac is contraindicated in patients in whom attacks of asthma, urticaria, or acute rhinitis have been observed following application of acetyl salicylic acid or other cyclo-oxygenase inhibitors.

## Nopain

### Sterile Eye Drops

**Precautions :** Patients tending toward uncontrolled bleeding or under therapy with anticoagulants should use Diclofenac with caution. If infection is present or there is a risk of infection, appropriate treatment (e.g. with antibiotics) should be given concomitantly.

**Side effects :** Occasional mild or moderate, transient stinging and/or blurred vision directly after application of eye drops. Hypersensitivity reaction in the form of itching, reddening, and photosensitivity; post-operative mydriasis have been reported rare.

**Pregnancy and lactation :** Pregnancy Category **C** by FDA.

**Packing :** 5ml in plastic dropper bottle.