

Composition : Omega-3-Acid Ethyl Esters 1gm (A combination of Eicosapentaenoic Acid & Docosahexaenoic Acid) Capsule.

Indication : **OMG-3** is indicated as an adjunct to diet to reduce very high (500mg/dL) triglyceride (TG) level in adult patients.

Dosage and administration : In Hypertriglyceridaemia, it can be taken as single dose 4 capsules (4gm) or 2 capsules (2gm) twice daily with food. Patients should be placed on an appropriate lipid lowering diet before receiving **OMG-3** and should continue this diet during treatment. Or, as directed by the registered physician.

Contraindication : Omega-3-acid ethyl esters is contraindicated in patients who exhibit hypersensitivity to any component of this medication.

Precautions : Omega-3-acid ethyl esters therapy should be withdrawn in patients who do not have an adequate response after 2 months of treatment.

Side effects : Adverse events are rarely reported such as enlarged abdomen, body odor, chest pain, chills, fever, generalized edema, hypertension, anorexia, constipation, drymouth, gastritis dizziness etc.

Use in pregnancy and lactation : There are no adequate and well-controlled studies in pregnant women.

OMG-3 Soft Capsule

Drug interactions : Patients receiving treatment with both Omega-3-acid ethyl esters and anticoagulants should be monitored periodically.

Packing : **OMG-3** : 6 x 5's soft capsules in blister pack.