

Composition : Paracetamol BP 125mg, 250mg & 500mg Suppository.

Pharmacology : Paracetamol is a non-opioid, non-salicylate analgesic. The site and mechanism for the analgesic effect of Paracetamol has not been determined but is thought to primarily involve central actions. Paracetamol is primarily metabolized in the liver by first-order kinetics and involves three principal separate pathways.

Indication : It is indicated for fever, common cold and influenza, headache, toothache, earache, bodyache, back pain, inflammatory pain, post-vaccination pain, fever of children & osteoarthritic pain.

Dosage & Administration : Suppository should be administered rectally.

> Children 3 months - 1 year: 1-2 suppository (60-120mg). The dosage should be based on age & weight i.e. 3 months (5kg)-60mg (1 suppository), 1 year (10kg)-120mg (2 suppositories of 60mg)

> Children below 5 years: 125-250mg, 2-3 times daily.

> Children 6-12 years: 250-500mg, 2-3 times daily.

Momodol
Suppository



> Adults & children over 12 years: 500-1000mg, 2-3 times daily.

Or, as directed by the registered physician.

Side effects : Side effects of paracetamol are usually mild, though haematological reactions including thrombocytopenia, leukopenia, pancytopenia, neutropenia has been reported. Pancreatitis, skin rashes and other allergic reaction occur occasionally.

Use in Pregnancy and Lactation : Paracetamol is safe in all stages of pregnancy and lactation

Overdose : There is no data available

Packing :

Momodol 125 Suppository : Each box contains 30 Suppositories in blister pack.

Momodol 250 Suppository : Each box contains 30 Suppositories in blister pack.

Momodol 500 Suppository : Each box contains 25 Suppositories in blister pack.