

Composition : Each ml eye drops contains Povidone BP 50 mg.

Pharmacology : Eyeteare Eye Drops contains Povidone BP 5%, a synthetic polymer consisting essentially of linear-1-Vinyl-2-Pyrrolidinone groups. Povidone is indicated for the management of dry eye as artificial tears.

Indications : Povidone Eye Drops is used for the symptomatic treatment of dry eye conditions including Keratoconjunctivitis sicca. It is also given as a substitute of tear fluid in case of unstable tear film or insufficient moistening of the eye surface.

Dosage & Administration : Adults (including the elderly): Instill 2 to 3 drops of the solution onto the eye/eyes and leave for two minutes or as required. Or, as directed by the registered physician.

How to use single dose :

- * Wash hands before start. Open plastic pouch & remove the plastic container.
- * Twist off the plastic containers cap.
- * Hold the plastic container pointing down, between thumb and fingers.
- * Tilt head back.
- * Pull down lower eyelid with a finger, until there is a 'pocket' between the eyelid and eye. The drop will go in here.
- * Bring the tip of the plastic container close to the eye. Do this in front of a mirror if it helps.
- * Do not touch eye or eyelid, surrounding areas or other surfaces with the plastic container.
- * Gently press on the base of the plastic container to release one drop at a time.

Contraindications : It is contraindicated in patients with known hypersensitivity to any ingredient of the product.

Precautions : Patients who experience blurred vision after application of the product should not drive or use machinery until their vision has

**Eyeteare
Eye Drops**



**DRUG
INTERNATIONAL
LTD.**

cleared. Contact lenses should not be worn during instillation of the product. After instillation there should be an interval of at least 30 minutes before reinsertion of contact lenses.

Side effects : Occasionally mild-transient burning or sticky sensation and very rarely irritation or hypersensitivity reactions are reported. Blurred vision after application may occur.

Use in Pregnancy and Lactation : Eyeteare 5% eye drops can be used during pregnancy. No effects on the breastfed/infant are anticipated since the systemic exposure of the breast-feeding woman to iodine is negligible.

Use in child : The adult dose may be used in infants, children and adolescents.

Drug- Interactions : In case of any additional local ocular treatment e.g. glaucoma therapy there should be an application interval of least 5 minutes between the two medications. Eyeteare always should be the last medication instilled.

Overdose : Not known and not likely about overdose.

Storage : Store below 30°C away from sunlight. Keep out of reach of children. Once the pouch has been opened, any unused plastic container should be stored in the open pouch. Dispose of the plastic container after use. Do not keep an open plastic container to use again.

Packing : Each box contains 25's (0.4ml) single use plastic containers enclosed in a plastic pouch overwrap.

